

RESEARCH NOTE

## Inhibitory Effects of Naringenin, Kaempferol, and Apigenin on Cholesterol Biosynthesis in HepG2 and MCF-7 Cells

Kee-Tae Kim<sup>1</sup>, Eun-Ju Yeo<sup>2</sup>, Sun-Hee Moon<sup>2</sup>, Ssang-Goo Cho<sup>1,2</sup>, Ye-Sun Han<sup>1,3</sup>, Seung-Yeol Nah<sup>1,4</sup>, and Hyun-Dong Paik<sup>1,2\*</sup>

<sup>1</sup>Bio/Molecular Informatics Center, Konkuk University, Seoul 143-701, Korea

<sup>2</sup>Division of Animal Life Science, Konkuk University, Seoul 143-701, Korea

<sup>3</sup>Department of Advanced Technology Fusion, Konkuk University, Seoul 143-701, Korea

<sup>4</sup>Department of Physiology, College of Veterinary Medicine, Konkuk University, Seoul 143-701, Korea

**Abstract** The inhibitory effects of naringenin, kaempferol, and apigenin on the production of cholesterol in HepG2 KCLB 88065 and MCF-7 KCLB 30022 cells were evaluated. In this study, quercetin was used as a reference reagent. After incubation for 3 days, fat-soluble contents of both cell types were extracted by using the Folch method and the cholesterol contents in both cultured cells were determined by high performance liquid chromatography. The concentration of cholesterol in untreated each tissue cells was  $12.2 \pm 0.11$  and  $8.83 \pm 0.12$  mg/g of lipid, respectively. The total concentration of each flavonoid was adjusted to 0, 35, or 350  $\mu$ M in the culture broth. As the results, the addition of 2% methanol and dimethyl sulfoxide (DMSO) to the media (control for flavonoid solvents) did not significantly affect cell growth; however, DMSO caused an increase in the production of cholesterol. Each flavonoid inhibited the production of cholesterol in both HepG2 and MCF-7 cells at the concentration of 35  $\mu$ M above. In addition, the inhibitory effect of kaempferol on the production of cholesterol in these cells was greater than the other flavonoids tested and HepG2 cells are more sensitive to flavonoids than MCF-7. From the results, the inhibitory effects of flavonoids on cholesterol production are different depending on the cell type.

**Keywords:** flavonoid, cholesterol biosynthesis, naringenin, kaempferol, apigenin, HepG2, MCF-7

### Introduction

Cholesterol is an indispensable component of many biological membranes, in which it regulates membrane fluidity and activates enzymes and transport mechanisms indirectly. However, excessive concentrations of cholesterol in the body result in the stifling of membrane and physiological functions and the development of atherosclerosis and emboli. That is, hypercholesterolemia is a major risk factor for cardiovascular diseases such as atherosclerosis, myocardial infarction, heart attacks, and cerebrovascular diseases, some of the leading causes of death in industrialized countries (1). Recently, it has been actively studied on lowering cholesterol by using probiotics (2-4).

Flavonoids comprise a large group of naturally occurring compounds widely distributed in the plant kingdom. Some of these compounds have been reported to contain various and potent biological activities including antioxidative, tissue-protective, and tumorigenic effects as well as the inhibition of hepatic cholesterol biosynthesis (5,6). In special, it has been known that quercetin has the inhibitory effect on cholesterol biosynthesis of HepG2. Therefore, the consumption of plant foods such as fruits, vegetables, red wines, and juices provides protection against various diseases, including cancer and cerebrovascular disease (7,8).

These functions have been demonstrated most convincingly using appropriate *in vitro* assays that provide insight into the underlying molecular mechanisms (5). In addition, the lipid accumulation process in animal cells has been thoroughly studied and it has been proposed that phenomenon results from scavenger receptor-mediated internalization of modified low-density lipoprotein (LDL) (9). This reinforces the idea that free radicals play an important role in the development of a variety of human diseases, particularly atherosclerosis. Animal studies have also been used to evaluate the inhibitory effects of flavonoids on cholesterol production, however the inhibitory mechanism is not known, possibly due to the presence of many other factors in animal tissues. However, it appeared that the inhibitory effects of flavonoids on cholesterol biosynthesis may be different according to tissue species (5).

Therefore, the purpose of this study is to evaluate naringenin, kaempferol, and apigenin (which are representative flavonoids in plants) as inhibitors of cholesterol production in HepG2 cell (a cancer cell of hepatocytes) and MCF-7 cell (a breast cancer cell) culture. This will provide important information relevant to the pharmaceutical industry with regard to hypercholesterolemia and hypertension.

### Materials and Methods


**Materials** HepG2 KCLB 88065 and MCF-7 KCLB 30022 cells were obtained from the Korean Cell Line Bank (KCLB; Seoul, Korea). Dulbecco's modified Engle's medium (DMEM) and Roxwell Park Memorial Institute (RPMI) 1640 medium was purchased from Gibco Lab.

\*Corresponding author: Tel: +82-2-2049-6011; Fax: +82-2-455-3082

E-mail: hdpaik@konkuk.ac.kr

Received April 21, 2008; Revised June 11, 2008;

Accepted June 22, 2008



**Fig. 1. Chemical structure of flavonoids.** (a) Naringenin, (b) kaempferol, (c) apigenin, and (d) quercetin.


(North Andover, MA, USA) and fetal bovine serum (FBS) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Both culture media were supplemented with 10% FBS and penicillin/streptomycin (100 U/mL each, Sigma-Aldrich). Naringenin, kaempferol, apigenin, and quercetin (Fig. 1) and cholesterol (free form, 5-cholesten-3 $\beta$ -ol) as a standard reagent for high performance liquid chromatography (HPLC) also were purchased from Sigma-Aldrich.

**Cell culture** HepG2 and MCF-7 cells were seeded in DMEM and RPMI containing various flavonoids. Flavonoids dissolved in 99% methanol or dimethyl sulfoxide (DMSO, Sigma-Aldrich) were diluted 1:50 in culture medium such that the total concentration of flavonoids was 0 (control), 35, or 350  $\mu$ M. Cultures were incubated in a humidified incubator (MCO-18AIC; Sanyo, Osaka, Japan) with a 5% CO<sub>2</sub> atmosphere at 37°C for 2 days (10-12).

**Extraction of crude cholesterol in cells** Lipids were recovered by using a modified method of Folch (13). That is, 100 mL of cell culture was added to 100 mL of solvent (chloroform:methanol=3:1), stirred rigorously, and extracted in a separatory funnel. The solvent layer was filtered and the filtrates were evaporated in a rotary vacuum evaporator (N-1000 series; Rikakikai Co., Tokyo, Japan). The concentrates were weighed, dissolved in methanol, and filtered with a 0.45- $\mu$ m membrane filter (MFS-25; Advantec MFS, Inc., Dublin, CA, USA) for further analysis.

**Cholesterol analysis** HPLC consisted of an Agilent 1100 pump and UV detector series (1100 series; Agilent Technologies, Inc., Palo Alto, CA, USA). Cholesterol as a standard reagent was purchased from Sigma-Aldrich. Twenty  $\mu$ L was injected into a  $\mu$ -Bondapak C<sub>18</sub> column (3.9 $\times$ 300 mm, Waters, Milford, Ireland) (14). The mobile phase was acetonitrile-methanol (3:1) and the flow rate was 1.0 mL/min. Detection was done at 202 nm.

**Statistical analysis** Analysis of variance was performed for each group of 3 samples using the SAS program. Duncan's test also was used to verify the significance of differences for each treatment.



**Fig. 2. HPLC chromatogram of cholesterol in HepG2 cells.**

## Results and Discussion

**Cholesterol contents of cultured cells** The HPLC pattern of cholesterol in HepG2 and MCF-7 cells is shown in Fig. 2. The concentration of cholesterol in untreated HepG2 and MCF-7 without the treatments was 12.2 $\pm$ 0.11 and 8.83 $\pm$ 0.12% in total lipid of cells, respectively (Table 1 and 2). In addition, the effects of 2.0% methanol and 2.0% DMSO itself in broth as controls for the flavonoid treatments on cholesterol biosynthesis were tested (DMSO also was used as a solvent because apigenin was insoluble in methanol at this concentration range). As the results, it appeared that both 2.0% methanol and 2.0% DMSO in the broth did not significantly affect the cell growth rate (data not shown). However, in case of cholesterol production in cells, 2.0% DMSO resulted in a slight increase in the production of cholesterol in cells, although the reason for these phenomena is still unknown.

### Effects of flavonoids on cholesterol production in cells

In the pharmaceutical industry, chemicals such as lovastatin and simvastatin have been used as inhibitors of cholesterol synthesis and to treat hypercholesterolemia (15) because these compounds inhibit 3-hydroxy-3-methylglutaryl CoA (HMG-CoA) reductase, which catalyzes the conversion of HMG-CoA to mevalonic acid as a rate-limiting step in cholesterol biosynthesis. In this pathway, it has been known that HMG-CoA reductase is regulated by phosphorylation/dephosphorylation (16).

Flavonoids as the natural materials also have been suggested to lower blood cholesterol levels by 2 kinds of mechanisms. Initially, it was suggested that flavonoids enhance the phosphorylation of HMG-CoA reductase indirectly and thus diminish endogenous cholesterol production (16). Secondly, flavonoids also probably exert their influence on steroid metabolism at other pivotal points. That is, flavonoids bind to the cytoplasmic steroid receptor due to the hydrophobicity of their aglycone portion, and this complex is likely to interact with steroid regulatory elements. Alternatively, the flavonoid may intercalate itself between the bases of this DNA segment and affect the transcription of genes involved in lowering the blood cholesterol level.

**Table 1. Effects of flavonoids in methanol on the cholesterol contents of HepG2 and MCF-7 cells**

Flavonoid	Conc. ( $\mu\text{M}$ )	Cholesterol content <sup>1)</sup> (%)	
		HepG2	MCF-7
Control	Media	12.2 $\pm$ 0.11	8.83 $\pm$ 0.12
	Methanol	11.5 $\pm$ 0.23	8.60 $\pm$ 0.05
Naringenin	35	0.34 $\pm$ 0.07	4.77 $\pm$ 0.11
	350	0.06 $\pm$ 0.02	4.06 $\pm$ 0.03
Kaempferol	35	2.27 $\pm$ 0.18	3.16 $\pm$ 0.20
	350	Trace	Trace
Quercetin <sup>2)</sup>	35	1.48 $\pm$ 0.23	7.35 $\pm$ 0.15
	350	0.52 $\pm$ 0.11	4.30 $\pm$ 0.23

<sup>1)</sup>mg/g of total lipid.<sup>2)</sup>Used as a reference reagent.**Table 2. Effects of flavonoids in DMSO on the cholesterol contents of HepG2 and MCF-7 cells**

Flavonoid	Conc. ( $\mu\text{M}$ )	Cholesterol content <sup>1)</sup> (%)	
		HepG2	MCF-7
Control	Medium	12.2 $\pm$ 0.11	8.83 $\pm$ 0.12
	DMSO	20.0 $\pm$ 0.23	15.4 $\pm$ 0.20
Naringenin	35	3.65 $\pm$ 0.15	4.15 $\pm$ 0.02
	350	2.50 $\pm$ 0.12	2.96 $\pm$ 0.13
Kaempferol	35	3.70 $\pm$ 0.04	4.26 $\pm$ 0.11
	350	Trace	Trace
Apigenin	35	9.18 $\pm$ 0.14	4.85 $\pm$ 0.20
	350	2.31 $\pm$ 0.09	0.77 $\pm$ 0.11
Quercetin <sup>2)</sup>	35	8.23 $\pm$ 0.02	6.77 $\pm$ 0.07
	350	Trace	3.16 $\pm$ 0.23

<sup>1)</sup>mg/g of total lipid.<sup>2)</sup>Used as a reference reagent.

In this study, quercetin was used as a reference because Gläßer *et al.* (5) explained that quercetin has inhibitory effects on cholesterol biosynthesis in HepG2 cells. According to the data in Table 1 and 2, it appeared that naringenin, kaempferol, apigenin, and quercetin affected the production of cholesterol in both HepG2 and MCF-7 cells at the concentration of 35  $\mu\text{M}$  above. As shown from Table 1, HepG2 cells were more sensitive to the inhibitory effects of flavonoids than MCF-7 cells. At a 35  $\mu\text{M}$  concentration in the methanol solvent in HepG2 cell cultures, naringenin was the most effective. However, kaempferol at a 350  $\mu\text{M}$  concentration was the most effective of the 4 flavonoids tested at lowering cholesterol in both tissues. In case of MCF-7 cells, kaempferol had the greatest inhibitory effects on cholesterol production in both methanol and DMSO solvents (Table 1 and 2). Overall, the flavonoids negatively affected the production of cholesterol in both cell lines at the concentration of 35  $\mu\text{M}$  above. These results support that the inhibitory effects of flavonoids toward the production of cholesterol vary depending on the flavonoids and the cell type. From these results, it appeared that the inhibitory effect of kaempferol on cholesterol production in cells is greater than that of the other

flavonoids tested.

Based on these studies, it appears that the inhibitory effects of flavonoids on the synthesis of cholesterol differ depending on the type and size of molecular groups and the positions of their side chains in the flavonoid backbone structure, and the cell type being treated. The detailed mechanisms and kinetics of the inhibitory effects of flavonoids are as yet unknown. However, the results of this study suggest that flavonoids can be used clinically to treat patients with hypercholesterolemia and hypertension. In future studies, the effective concentration and synergistic effects of flavonoids used in combination will be determined.

## Acknowledgments

This work was supported by the KRF 2006-005-J03403 and the Brain Korea 21 program, Korea.

## References

- Wald NJ, Law MR. Serum cholesterol and ischemic heart disease. *Atherosclerosis* 118: S1-S5 (1995)
- Ha CG, Cho JK, Lee CH, Chai YC, Ha YA, Shin SH. Cholesterol lowering effect of *Lactobacillus plantarum* isolated from human feces. *J. Microbiol. Biotechnol.* 16: 1201-1209 (2006)
- Lee HW, Chien JT, Chen BH. Inhibition of cholesterol oxidation in marinated foods as affected by antioxidants during heating. *Food Chem.* 108: 234-244 (2008)
- Chien JT, Hsu DJ, Chen BH. Kinetic model for studying the effect of quercetin on cholesterol oxidation during heating. *J. Agr. Food Chem.* 54: 1486-1492 (2006)
- Gläßer G, Graefe EU, Struck F, Veit M, Gebhardt R. Comparison of antioxidative capacities and inhibitory effects on cholesterol biosynthesis of quercetin and potential metabolites. *Phytomedicine* 9: 33-40 (2002)
- Lee JY, Moon SO, Kwon YJ, Rhee SJ, Park HR, Choi SW. Identification and quantification of anthocyanins and flavonoids in mulberry cultivars. *Food Sci. Biotechnol.* 13: 176-184 (2004)
- Choi YM, Ku JB, Chang HB, Lee J. Antioxidant activities and total phenolics of ethanol extracts from several edible mushrooms produced in Korea. *Food Sci. Biotechnol.* 14: 700-703 (2005)
- Morré DM, Morré DJ. Anticancer activity of grape and grape skin extracts alone and combined with green tea infusions. *Cancer Lett.* 238: 202-209 (2006)
- Laurence G, Nadine L, Denis B. Role of the cyclic AMP-dependent pathway in free radical-induced cholesterol accumulation in vascular smooth muscle cells. *Free Radical Bio. Med.* 29: 181-190 (2000)
- Funatsu T, Suzuki K, Goto M, Arai Y, Kakuta H, Tanaka H, Yasuda S, Ida M, Nishijima S, Miyata K. Prolonged inhibition of cholesterol synthesis by atorvastatin inhibits apoB-100 and triglyceride secretion from HepG2 cells. *Atherosclerosis* 157: 107-115 (2001)
- Miki I, Teruo M. The growth inhibitory effect of conjugated linoleic acid on a human hepatoma cell line, HepG2, is induced by a change in fatty acid metabolism, but not the facilitation of lipid peroxidation in the cells. *Biochim. Biophys. Acta* 1530: 162-171 (2001)
- Lee YH, Ho JN, Dong MS, Park CH, Kim HK, Hong BS, Shin DH, Cho HY. Transfected HepG2 cells for evaluation of catechin effects on alcohol-induced CYP2E1 cytotoxicity. *J. Microbiol. Biotechnol.* 15: 1310-1316 (2005)
- Suleen SH, Sebely P. Margarine phytosterols decrease the secretion of atherogenic lipoproteins from HepG2 liver and Caco2 intestinal cells. *Atherosclerosis* 182: 29-36 (2005)
- Eugenios K, Pau. BA. Novel HPLC analysis of tocopherols, tocotrienols, and cholesterol in tissue. *Free Radical Bio. Med.* 27: 1137-1140 (1999)
- Pazzucconi F, Dorigotti F, Gianfranceschi G, Campagnoli G

- Champagnoli G, Sirtori M, Franceschini G, Sirtori CR. Therapy with HMG CoA reductase inhibitors: Characteristics of the long-term permanence of hypocholesterolemic activity. *Atherosclerosis* 117: 189-198 (1995)
16. Stein JH, Keevil JG, Wiebe DA, Aaschlimann S, Fols JD. Purple grape juice improves endothelial function and reduces the susceptibility of LDL cholesterol to oxidation in patients with coronary disease. *Circulation* 100: 1050-1055 (1999)